

in a sucrose molecule to form raffinose, wherein said polynucleotide comprises a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence encoding the amino acid sequence as depicted in SEQ ID NO: 1,

(b) a nucleotide sequence as depicted in SEQ ID NO: 2,

(c) a nucleotide sequence encoding the amino acid sequence as depicted in SEQ ID NO: 3,

(d) a nucleotide sequence depicted by the 236th to 2584th nucleotides in the nucleotide sequence as depicted in SEQ ID NO: 4,

FI
cont (e) a nucleotide sequence encoding the amino acid sequence as depicted in SEQ ID NO: 5,

(f) a nucleotide sequence depicted by the 134th to 2467th nucleotides in the nucleotide sequence as depicted in SEQ ID NO: 6,

(g) a nucleotide sequence encoding the amino acid sequence as depicted in SEQ ID NO: 7,

(h) a nucleotide sequence depicted by the 1st to 1719th nucleotides in the nucleotide sequence as depicted in SEQ ID NO: 8,

(i) a nucleotide sequence obtainable from a polynucleotide which is amplifiable from a nucleic acid obtained from soybean with

a combination of a PCR primer of SEQ ID NO: 9 and a PCR primer of SEQ ID NO: 10, wherein said nucleotide sequence is hybridizable with a nucleotide sequence complementary to the nucleotide sequence of (a) or (b), in a buffer comprising 0.9M NaCl and 0.09M citric acid at 65°C to 68°C,

FI
cont
(j) a nucleotide sequence obtainable from a polynucleotide which is amplifiable from a nucleic acid obtained from a *Chenopdiaceae* plant with a combination of a PCR primer selected from the group consisting of SEQ ID NO: 11 and SEQ ID NO: 13 and a PCR primer selected from the group consisting of SEQ ID NO: 12 and SEQ ID NO: 14, wherein said nucleotide sequence is hybridizable with a nucleotide sequence complementary to the nucleotide sequence of (c) or (d), in a buffer comprising 0.9M NaCl and 0.09M citric acid at 65°C to 68°C, and

(k) a nucleotide sequence obtainable from a polynucleotide which is amplifiable from a nucleic acid obtained from a *Cruciferae* plant with a combination of a PCR primer selected from the group consisting of SEQ ID NO: 15, SEQ ID NO: 17 and SEQ ID NO: 19 and a PCR primer selected from the group consisting of SEQ ID NO: 16, SEQ ID NO: 18 and SEQ ID NO: 20, wherein said nucleotide sequence is hybridizable with a nucleotide sequence complementary to the nucleotide sequence of any one of (e) to (h), in a buffer comprising 0.9M NaCl and 0.09M citric acid at 65°C to 68°C.

30. (Twice Amended) An isolated nucleic acid which comprises a polynucleotide encoding a protein that binds a D-galactosyl group through the $\alpha(1\rightarrow6)$ bond to the hydroxyl group attached to the carbon atom at 6-position of the D-glucose residue in a sucrose molecule to form raffinose, wherein said polynucleotide comprises a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence encoding the amino acid sequence as depicted in SEQ ID NO: 1,

(b) a nucleotide sequence as depicted in SEQ ID NO: 2,

fr (c) a nucleotide sequence encoding the amino acid sequence as depicted in SEQ ID NO: 3,

(d) a nucleotide sequence depicted by the 236th to 2584th nucleotides in the nucleotide sequence as depicted in SEQ ID NO: 4,

(e) a nucleotide sequence encoding the amino acid sequence as depicted in SEQ ID NO: 5,

(f) a nucleotide sequence depicted by the 134th to 2467th nucleotides in the nucleotide sequence as depicted in SEQ ID NO: 6,

(g) a nucleotide sequence encoding the amino acid sequence as depicted in SEQ ID NO: 7,

(h) a nucleotide sequence depicted by the 1st to 1719th nucleotides in the nucleotide sequence as depicted in SEQ ID NO: 8,

(i) a nucleotide sequence obtainable from a polynucleotide which is amplifiable from a nucleic acid obtained from soybean with a combination of a PCR primer of SEQ ID NO: 9 and a PCR primer of SEQ ID NO: 10, wherein said nucleotide sequence is hybridizable with a nucleotide sequence complementary to the nucleotide sequence of (a) or (b), in a buffer comprising 0.9M NaCl and 0.09M citric acid at 65°C to 68°C,

F2
cond
(j) a nucleotide sequence obtainable from a polynucleotide which is amplifiable from a nucleic acid obtained from beet with a combination of a PCR primer selected from the group consisting of SEQ ID NO: 11 and SEQ ID NO: 13 and a PCR primer selected from the group consisting of SEQ ID NO: 12 and SEQ ID NO: 14, wherein said nucleotide sequence is hybridizable with a nucleotide sequence complementary to the nucleotide sequence of (c) or (d), in a buffer comprising 0.9M NaCl and 0.09M citric acid at 65°C to 68°C, and

(k) a nucleotide sequence obtainable from a polynucleotide which is amplifiable from a nucleic acid obtained from a *Cruciferae* plant with a combination of a PCR primer selected from the group consisting of SEQ ID NO: 15, SEQ ID NO: 17 and SEQ ID NO: 19 and a PCR primer selected from the group consisting of SEQ ID NO: 16, SEQ ID NO: 18 and SEQ ID NO: 20, wherein said nucleotide sequence is

hybridizable with a nucleotide sequence complementary to the
F2
cond nucleotide sequence of any one of (e) to (h), in a buffer
comprising 0.9M NaCl and 0.09M citric acid at 65°C to 68°C.

Attached hereto is a marked up version showing the changes
made to the application by this Amendment.